

AMANITA MUSCARIA



Psychoactive Compounds

A concise guide to muscimol, ibotenic acid, brain mechanisms, subjective effects, and toxicology

EIRLAB Scientific overview · October 2026

CONTENTS

1. Introduction
2. The two principal psychoactive compounds
3. The subjective experience
4. Pharmacological comparison
5. Duration and time course
6. Preparation and chemical conversion
7. Comparison with psilocybin mushrooms
8. Risks and harm reduction
9. Key conclusions
10. Further reading

01 — INTRODUCTION

Amanita muscaria, commonly known as fly agaric, is a distinctive red-capped mushroom with white spots. Unlike psilocybin-containing mushrooms, its principal psychoactive compounds act primarily on the brain's GABA and glutamate systems.

Its effects can include sedation, dreamlike perception, altered awareness, impaired coordination, agitation, and delirium-like confusion.

The experience is highly variable because the concentrations of active compounds differ between individual mushrooms and preparations. The mushroom should therefore be understood as a potentially toxic, neuroactive organism rather than a predictable psychedelic.

02 — THE TWO PRINCIPAL COMPOUNDS

Muscimol

Primary mechanism: GABA-A receptor agonism.

Muscimol activates GABA-A receptors, which mediate much of the brain's inhibitory neurotransmission.

Potential effects

- Drowsiness and sedation
- Dreamlike states and altered awareness
- Distorted perception and time perception
- Impaired balance and coordination
- Confusion and reduced responsiveness at higher exposures

Muscimol is considered a major driver of the characteristic intoxicating effects of fly agaric. Its effects are not limited to sleepiness: interference with neural networks involved in perception, movement, and consciousness can produce unusual and disorienting experiences.

Ibotenic acid

Primary mechanism: Agonism at glutamate receptors, including NMDA receptors.

Ibotenic acid activates excitatory neurotransmitter pathways. It may contribute to perceptual disruption, agitation, involuntary movements, and neurological toxicity.

It is also a chemical precursor to muscimol. Through a reaction called decarboxylation, ibotenic acid can lose carbon dioxide and form muscimol.

This conversion can occur during drying or heating, but the extent depends on the conditions. It cannot be used as a reliable indicator that a preparation is safe.

The essential distinction

Muscimol primarily activates inhibitory signalling, whereas ibotenic acid activates excitatory signalling. These effects do not simply cancel each other out. Together, the compounds can disrupt normal brain function in unpredictable ways.

03 — WHAT THE EXPERIENCE CAN FEEL LIKE

Reported experiences vary considerably. The following are broad patterns, not guaranteed effects.

Dreamlike perception

The environment may seem unreal, distant, distorted, or dreamlike. Some people report changes in time perception, visual distortions, or experiences resembling the transition between waking and sleep.

Sedation and reduced alertness

Muscimol can produce physical heaviness, drowsiness, reduced alertness, and unusual dreamlike states. More severe intoxication may result in profound sedation or loss of consciousness.

Agitation and confusion

Ibotenic acid's excitatory activity may contribute to restlessness, disorientation, agitation, and involuntary movements. A person may alternate between apparent stimulation and sleepiness.

Motor impairment

Unsteadiness, stumbling, dizziness, slurred speech, and poor judgement can occur. Significant impairment may be present even when the person subjectively feels comfortable.

Physical symptoms

Nausea, vomiting, muscle twitching, changes in heart rate, and temperature disturbances are possible. Severe cases can involve seizures or coma.

The contribution of each compound to an individual's symptoms cannot always be determined from the clinical presentation alone.

04 — PHARMACOLOGICAL COMPARISON

Feature	Muscimol	Ibotenic acid
Main receptor action	GABA-A agonist	Glutamate-receptor agonist
Broad neural effect	Inhibitory signalling	Excitatory signalling
Typical effects	Sedation, dreamlike states	Agitation, perceptual disruption

Possible motor effects	Unsteadiness, poor coordination	Involuntary movements, poor coordination
Serious risks	Reduced consciousness	Severe neurological disturbance

These are broad pharmacological tendencies. They do not describe two separate experiences that can reliably be distinguished during intoxication.

Why muscimol is not simply a sedative

GABA-A receptors occur throughout the brain and influence multiple neural circuits. Activating them can affect perception, memory, movement, and consciousness as well as alertness.

Consequently, muscimol intoxication may involve dreamlike hallucinations, disorientation, or unusual behaviour rather than ordinary sleepiness alone.

Why ibotenic acid warrants caution

Glutamate signalling is essential for normal brain function, but excessive activation of certain glutamate pathways can damage neurons under experimental conditions.

Ibotenic acid is used in neuroscience research to produce targeted brain lesions when administered directly into specific brain regions. This does not mean every oral exposure causes permanent brain damage; it does illustrate why the compound should not be regarded as a benign stimulant.

05 — DURATION AND TIME COURSE

Effects are commonly reported to begin within approximately 30 minutes to two hours after ingestion, although onset varies. Intoxication may intensify over the following hours and can last several hours. Severe cases may persist considerably longer.

Onset: Changes in alertness, perception, balance, or behaviour may begin.

Intensification: Sedation, confusion, dreamlike experiences, agitation, or involuntary movements may become more pronounced.

Resolution: Symptoms generally subside over time, although fatigue, confusion, or impaired coordination may linger.

These are approximate clinical patterns, not a dependable timetable. Delayed onset does not indicate that a preparation is inactive, and taking more while waiting for effects can increase the risk of severe intoxication.

06 — PREPARATION AND CHEMICAL CONVERSION

Ibotenic acid can be converted into muscimol through decarboxylation.

IBOTENIC ACID → MUSCIMOL

This chemical transformation can alter the relative concentrations of the two psychoactive compounds.

Important considerations:

- Drying, heating, and storage can change compound concentrations.
- The extent of conversion depends on the conditions.
- Boiling or soaking may remove some water-soluble compounds, but does not guarantee that toxins have been eliminated.
- Mushroom appearance and weight cannot reliably establish the concentration of active compounds.
- A preparation containing more muscimol is not automatically safe.

There is insufficient evidence to establish a reliable preparation method that consistently produces a safe recreational product.

07 — COMPARISON WITH PSILOCYBIN MUSHROOMS

Feature	<i>Amanita muscaria</i>	Psilocybin mushrooms
Principal psychoactive compounds	Muscimol and ibotenic acid	Psilocybin, converted to psilocin
Main mechanism	GABA and glutamate systems	Primarily serotonin 5-HT2A receptors
Characteristic effects	Dreamlike, sedating, disorienting	Altered perception, thought, and emotion
Coordination	Frequently impaired	Can also be impaired
Important risks	Confusion, fluctuating consciousness, neurological toxicity	Anxiety, panic, psychological distress, and other acute effects

Neither category is inherently safe. The key distinction is pharmacological: fly agaric produces a different form of intoxication, with potentially pronounced fluctuations in alertness and behaviour.

08 — RISKS AND HARM REDUCTION

The desired subjective effect cannot reliably be separated from the risk of toxicity. An apparently pleasant experience may coexist with substantial impairment.

Important risks include:

- Severe confusion, agitation, panic, or delirium
- Falls, accidents, and impaired judgement
- Vomiting and dehydration
- Seizures or profound sedation in severe cases

- Dangerous interactions with alcohol, sedatives, sleeping medications, and other psychoactive substances

Evidence on human dose-response relationships, long-term safety, and therapeutic benefits remains limited.

Emergency response

If someone has consumed fly agaric and develops severe confusion, seizures, breathing difficulties, or reduced consciousness, seek emergency medical help immediately.

Do not induce vomiting or leave the person alone. In Ireland, call **112 or 999** for a medical emergency.

09 — KEY CONCLUSIONS

Muscimol activates GABA-A receptors and contributes to sedation, altered consciousness, and dreamlike effects.

Ibotenic acid activates glutamate receptors and can contribute to excitation, perceptual disruption, and neurological toxicity.

Together, the compounds can produce a fluctuating mixture of sedation, altered perception, confusion, and motor impairment.

From a recreational perspective, *Amanita muscaria* is best understood as a toxic mushroom with an unpredictable intoxication profile, not as a predictable substitute for conventional psychedelics.

Its pharmacology is scientifically interesting, but uncertainty around composition, preparation, and human effects remains a substantial concern.



IRISH CANNABIS WHOLESALE